

An upper bound on the fractional chromatic number of triangle-free subcubic graphs

Chun-Hung Liu

Georgia Institute of Technology

May 10, 2012

Definition

An $(a : b)$ -coloring of a graph G is a function f which maps the vertices of G into b -element subsets of some set of size a in such a way that $f(u)$ is disjoint from $f(v)$ for every two adjacent vertices u and v in G .

Definition

An $(a : b)$ -coloring of a graph G is a function f which maps the vertices of G into b -element subsets of some set of size a in such a way that $f(u)$ is disjoint from $f(v)$ for every two adjacent vertices u and v in G .

Every $(a : 1)$ -coloring is a proper vertex a -coloring.

Definition

An $(a : b)$ -coloring of a graph G is a function f which maps the vertices of G into b -element subsets of some set of size a in such a way that $f(u)$ is disjoint from $f(v)$ for every two adjacent vertices u and v in G .

Every $(a : 1)$ -coloring is a proper vertex a -coloring.

The *fractional chromatic number* $\chi_f(G)$ is the infimum of a/b over all pairs of positive integers a, b such that G has an $(a : b)$ -coloring.

Definition

An $(a : b)$ -coloring of a graph G is a function f which maps the vertices of G into b -element subsets of some set of size a in such a way that $f(u)$ is disjoint from $f(v)$ for every two adjacent vertices u and v in G .

Every $(a : 1)$ -coloring is a proper vertex a -coloring.

The *fractional chromatic number* $\chi_f(G)$ is the infimum of a/b over all pairs of positive integers a, b such that G has an $(a : b)$ -coloring. So $\chi_f(G) \leq \chi(G)$.

Definition

An $(a : b)$ -coloring of a graph G is a function f which maps the vertices of G into b -element subsets of some set of size a in such a way that $f(u)$ is disjoint from $f(v)$ for every two adjacent vertices u and v in G .

Every $(a : 1)$ -coloring is a proper vertex a -coloring.

The *fractional chromatic number* $\chi_f(G)$ is the infimum of a/b over all pairs of positive integers a, b such that G has an $(a : b)$ -coloring. So $\chi_f(G) \leq \chi(G)$.

In fact, the infimum is the minimum.

A conjecture

If G has an $(a : b)$ -coloring, then $\alpha(G) \geq bn/a$.

A conjecture

If G has an $(a : b)$ -coloring, then $\alpha(G) \geq bn/a$.

Staton (1979) proved that $\alpha(G) \geq 5n/14$ for every triangle-free subcubic graph (i.e. graph of max. degree ≤ 3).

A conjecture

If G has an $(a : b)$ -coloring, then $\alpha(G) \geq bn/a$.

Staton (1979) proved that $\alpha(G) \geq 5n/14$ for every triangle-free subcubic graph (i.e. graph of max. degree ≤ 3).

Fajtlowicz (1978) pointed out that the generalized Petersen graph $P(7, 2)$ has 14 vertices and no independent set of size 6.

A conjecture

If G has an $(a : b)$ -coloring, then $\alpha(G) \geq bn/a$.

Staton (1979) proved that $\alpha(G) \geq 5n/14$ for every triangle-free subcubic graph (i.e. graph of max. degree ≤ 3).

Fajtlowicz (1978) pointed out that the generalized Petersen graph $P(7, 2)$ has 14 vertices and no independent set of size 6.

Heckman and Thomas (2001) gave a short proof of Staton's Theorem, and they gave the following conjecture.

Conjecture: *The fractional chromatic number of every triangle-free subcubic graph is at most $14/5$.*

A conjecture

If G has an $(a : b)$ -coloring, then $\alpha(G) \geq bn/a$.

Staton (1979) proved that $\alpha(G) \geq 5n/14$ for every triangle-free subcubic graph (i.e. graph of max. degree ≤ 3).

Fajtlowicz (1978) pointed out that the generalized Petersen graph $P(7, 2)$ has 14 vertices and no independent set of size 6.

Heckman and Thomas (2001) gave a short proof of Staton's Theorem, and they gave the following conjecture.

Conjecture: *The fractional chromatic number of every triangle-free subcubic graph is at most $14/5$.*

The statement that $\chi_f(G) \leq 14/5$ is equivalent to the statement that for every weight function defined on $V(G)$, there is an independent set I of G such that the sum of weights of vertices in I is at least $5w(G)/14$.

Known results

Let G be a triangle-free subcubic graph.

- ① (Hatami and Zhu 2009) $\chi_f(G) \leq 3 - \frac{3}{64} \approx 2.953$, and $\chi_f(G) \leq 2.78571$ if the girth of G is at least 7.

Known results

Let G be a triangle-free subcubic graph.

- ① (Hatami and Zhu 2009) $\chi_f(G) \leq 3 - \frac{3}{64} \approx 2.953$, and $\chi_f(G) \leq 2.78571$ if the girth of G is at least 7.
- ② (Lu and Peng 2010⁺) $\chi_f(G) \leq 3 - \frac{3}{43} \approx 2.930$.

Known results

Let G be a triangle-free subcubic graph.

- ① (Hatami and Zhu 2009) $\chi_f(G) \leq 3 - \frac{3}{64} \approx 2.953$, and $\chi_f(G) \leq 2.78571$ if the girth of G is at least 7.
- ② (Lu and Peng 2010⁺) $\chi_f(G) \leq 3 - \frac{3}{43} \approx 2.930$.
- ③ (Ferguson, Kaiser, and Král 2012⁺) $\chi_f(G) \leq 32/11 \approx 2.909$.

Known results

Let G be a triangle-free subcubic graph.

- 1 (Hatami and Zhu 2009) $\chi_f(G) \leq 3 - \frac{3}{64} \approx 2.953$, and $\chi_f(G) \leq 2.78571$ if the girth of G is at least 7.
- 2 (Lu and Peng 2010⁺) $\chi_f(G) \leq 3 - \frac{3}{43} \approx 2.930$.
- 3 (Ferguson, Kaiser, and Král 2012⁺) $\chi_f(G) \leq 32/11 \approx 2.909$.

Theorem (L.)

The fractional chromatic number of every triangle-free subcubic graph is at most $43/15 \approx 2.867$.

Larger maximum degree

Let G be a K_Δ -free graph of maximum degree Δ other than C_8^2 and $C_5 \boxtimes K_2$.

- ① (King, Lu, and Peng 2012) $\chi_f(G) \leq 4 - \frac{2}{67} \approx 3.9701$ when $\Delta = 4$. ($\chi_f(C_{11}^2) = 4 - \frac{1}{3} \approx 3.6778$).
- ② (King, Lu, and Peng 2012) $\chi_f(G) \leq 5 - \frac{2}{67} \approx 4.9701$ when $\Delta = 5$. ($\chi_f(C_7 \boxtimes K_2) = 5 - \frac{1}{3} \approx 4.6778$).
- ③ (Edwards and King) $\chi_f(G) \leq 6 - \frac{153}{3431} \approx 5.9554$ when $\Delta = 6$. ($\chi_f((C_5 \boxtimes K_3) - 4v) = 6 - \frac{1}{2} = 5.5$).
- ④ (Edwards and King) $\chi_f(G) \leq 7 - \frac{80}{889} \approx 6.9100$ when $\Delta = 7$. ($\chi_f((C_5 \boxtimes K_3) - 2v) = 7 - \frac{1}{2} = 6.5$).
- ⑤ (Edwards and King) $\chi_f(G) \leq 8 - \frac{17280}{152209} \approx 7.8864$ when $\Delta = 8$. ($\chi_f(C_5 \boxtimes K_3) = 8 - \frac{1}{2} = 7.5$).
- ⑥ (Edwards and King) $\chi_f(G) \leq 9 - \frac{17}{130} \approx 8.8692$ when $\Delta = 9$.
- ⑦ (Edwards and King) $\chi_f(G) \leq 10 - \frac{8565625}{60177971} \approx 9.8577$ when $\Delta = 10$.

Main ideas

Find a "good" proper 3-coloring f of G such that for each $1 \leq i < j \leq 3$, the subgraph, denoted by $G^{(i,j)}$, of G induced by vertices of color i and j has some "good" structures.

Main ideas

Find a "good" proper 3-coloring f of G such that for each $1 \leq i < j \leq 3$, the subgraph, denoted by $G^{(i,j)}$, of G induced by vertices of color i and j has some "good" structures.

Define two maps $g_1^{(i,j)}$ and $g_2^{(i,j)}$ from $V(G)$ to subsets of $[14]$ such that adjacent vertices receive disjoint sets, and

$$\begin{aligned} & |g_1^{(i,j)}(v)| + |g_2^{(i,j)}(v)| \\ &= \begin{cases} 4, & \text{if } f(v) \notin \{i, j\}; \\ 16 - 2\deg_{G^{(i,j)}}(v) - 2 \cdot 1_{I^{(i,j)}}(v), & \text{if } f(v) \in \{i, j\}. \end{cases} \end{aligned}$$

for some independent set $I^{(i,j)} \subseteq \{v : \deg_{G^{(i,j)}}(v) = 3\}$ of G .

Main ideas

Define $g^{(i,j)}$ by the map from $V(G)$ to subsets of $[28]$ such that $g^{(i,j)}(v) = g_1^{(i,j)}(v) \cup (g_2^{(i,j)}(v) + 14)$ for every $1 \leq i < j \leq 3$ and vertex v .

Main ideas

Define $g^{(i,j)}$ by the map from $V(G)$ to subsets of $[28]$ such that $g^{(i,j)}(v) = g_1^{(i,j)}(v) \cup (g_2^{(i,j)}(v) + 14)$ for every $1 \leq i < j \leq 3$ and vertex v .

Define g by the map from $V(G)$ to subsets of $[84]$ such that $g(v) = g^{(1,2)}(v) \cup (g^{(1,3)} + 28) \cup (g^{(2,3)}(v) + 56)$ for every vertex v .

Main ideas

Define $g^{(i,j)}$ by the map from $V(G)$ to subsets of $[28]$ such that $g^{(i,j)}(v) = g_1^{(i,j)}(v) \cup (g_2^{(i,j)}(v) + 14)$ for every $1 \leq i < j \leq 3$ and vertex v .

Define g by the map from $V(G)$ to subsets of $[84]$ such that $g(v) = g^{(1,2)}(v) \cup (g^{(1,3)} + 28) \cup (g^{(2,3)}(v) + 56)$ for every vertex v . So for vertex v with $f(v) = 1$,

$$\begin{aligned} |g(v)| &\geq 4 + 32 - 2(\deg_{G^{(1,2)}}(v) + \deg_{G^{(1,3)}}(v)) - 2(1_{I^{(1,2)}}(v) + 1_{I^{(1,3)}}(v)) \\ &= 36 - 2\deg_G(v) - 2(1_{I^{(1,2)}}(v) + 1_{I^{(1,3)}}(v) + 1_{I^{(2,3)}}(v)) \\ &\geq 30 - 2(1_{I^{(1,2)}}(v) + 1_{I^{(1,3)}}(v) + 1_{I^{(2,3)}}(v)). \end{aligned}$$

Main ideas

Define $g^{(i,j)}$ by the map from $V(G)$ to subsets of $[28]$ such that $g^{(i,j)}(v) = g_1^{(i,j)}(v) \cup (g_2^{(i,j)}(v) + 14)$ for every $1 \leq i < j \leq 3$ and vertex v .

Define g by the map from $V(G)$ to subsets of $[84]$ such that $g(v) = g^{(1,2)}(v) \cup (g^{(1,3)}(v) + 28) \cup (g^{(2,3)}(v) + 56)$ for every vertex v . So for vertex v with $f(v) = 1$,

$$\begin{aligned} |g(v)| &\geq 4 + 32 - 2(\deg_{G^{(1,2)}}(v) + \deg_{G^{(1,3)}}(v)) - 2(1_{I^{(1,2)}}(v) + 1_{I^{(1,3)}}(v)) \\ &= 36 - 2\deg_G(v) - 2(1_{I^{(1,2)}}(v) + 1_{I^{(1,3)}}(v) + 1_{I^{(2,3)}}(v)) \\ &\geq 30 - 2(1_{I^{(1,2)}}(v) + 1_{I^{(1,3)}}(v) + 1_{I^{(2,3)}}(v)). \end{aligned}$$

Similarly, $|g(v)| \geq 30 - 2(1_{I^{(1,2)}}(v) + 1_{I^{(1,3)}}(v) + 1_{I^{(2,3)}}(v))$ for every vertex v . So g is "almost" a $(84 : 30)$ -coloring.

Main ideas

Define $g^{(i,j)}$ by the map from $V(G)$ to subsets of $[28]$ such that $g^{(i,j)}(v) = g_1^{(i,j)}(v) \cup (g_2^{(i,j)}(v) + 14)$ for every $1 \leq i < j \leq 3$ and vertex v .

Define g by the map from $V(G)$ to subsets of $[84]$ such that $g(v) = g^{(1,2)}(v) \cup (g^{(1,3)} + 28) \cup (g^{(2,3)}(v) + 56)$ for every vertex v . So for vertex v with $f(v) = 1$,

$$\begin{aligned} |g(v)| &\geq 4 + 32 - 2(\deg_{G^{(1,2)}}(v) + \deg_{G^{(1,3)}}(v)) - 2(1_{I^{(1,2)}}(v) + 1_{I^{(1,3)}}(v)) \\ &= 36 - 2\deg_G(v) - 2(1_{I^{(1,2)}}(v) + 1_{I^{(1,3)}}(v) + 1_{I^{(2,3)}}(v)) \\ &\geq 30 - 2(1_{I^{(1,2)}}(v) + 1_{I^{(1,3)}}(v) + 1_{I^{(2,3)}}(v)). \end{aligned}$$

Similarly, $|g(v)| \geq 30 - 2(1_{I^{(1,2)}}(v) + 1_{I^{(1,3)}}(v) + 1_{I^{(2,3)}}(v))$ for every vertex v . So g is "almost" a $(84 : 30)$ -coloring.

A minor modification yields an $(86 : 30)$ -coloring.

Main ideas

$I^{(1,2)}$, $I^{(1,3)}$, and $I^{(2,3)}$ are pairwise disjoint independent sets, and the union of them is also an independent set.

Main ideas

$I^{(1,2)}$, $I^{(1,3)}$, and $I^{(2,3)}$ are pairwise disjoint independent sets, and the union of them is also an independent set.



Let $I = I^{(1,2)} \cup I^{(1,3)} \cup I^{(2,3)}$.

Define g' by $g'(v) = g(v)$ if $v \notin I$, and $g'(v) \cup \{85, 86\}$ if $v \in I$.

Hence, g' is a $(86 : 30)$ -coloring, so $\chi_f(G) \leq 86/30 = 43/15$.

Graph L_0

Let L_0 be the graph that is obtained by identifying two paths of order four of two different C_5 's. A L_0 is a *rainbow* L_0 with respect to a proper 3-coloring f if the shared 4-path is bicolored by f .



Good 3-coloring


A proper 3-coloring f of a subcubic graph G is *good* if for every $1 \leq i < j \leq 3$:

- G contains no rainbow L_0 's with respect to f .


Good 3-coloring

A proper 3-coloring f of a subcubic graph G is *good* if for every $1 \leq i < j \leq 3$:


- G contains no rainbow L_0 's with respect to f .



Good 3-coloring



Good 3-coloring



Fractionally critical graph

G is *fractionally t -critical* if $\chi_f(G) > t$ but $\chi_f(H) \leq t$ for every proper subgraph H of G .

Lemma

If G is a fractionally t -critical triangle-free subcubic graph, where $t \geq 8/3$, then G has a good 3-coloring.

Next step

We have a good 3-coloring h , and now we want to define $g_1^{(i,j)}$ and $g_2^{(i,j)}$ from $V(G)$ to subsets of $[14]$ such that adjacent vertices receive disjoint sets, and

$$\begin{aligned} & |g_1^{(i,j)}(v)| + |g_2^{(i,j)}(v)| \\ &= \begin{cases} 4, & \text{if } h(v) \notin \{i, j\}; \\ 16 - 2\deg_{G^{(i,j)}}(v) - 2 \cdot 1_{I^{(i,j)}}(v), & \text{if } h(v) \in \{i, j\}. \end{cases} \end{aligned}$$

Next step

We have a good 3-coloring h , and now we want to define $g_1^{(i,j)}$ and $g_2^{(i,j)}$ from $V(G)$ to subsets of $[14]$ such that adjacent vertices receive disjoint sets, and

$$\begin{aligned} & |g_1^{(i,j)}(v)| + |g_2^{(i,j)}(v)| \\ &= \begin{cases} 4, & \text{if } h(v) \notin \{i, j\}; \\ 16 - 2\deg_{G^{(i,j)}}(v) - 2 \cdot 1_{I^{(i,j)}}(v), & \text{if } h(v) \in \{i, j\}. \end{cases} \end{aligned}$$

We first define each $g_1^{(i,j)}(v)$ and $g_2^{(i,j)}(v)$ by 2 colors on vertices v with $h(v) \notin \{i, j\}$, and then extend the coloring to $G^{(2,3)}$.

Remark

We can prove that $\chi_f(G) \leq 14/5$ if one can show that every fractionally $14/5$ -critical triangle-free subcubic graph has a good 3-coloring such that each two color classes induced a subgraph of maximum degree at most two.

THANK YOU

HAPPY BIRTHDAY, ROBIN!